



Product information

Lebosol[®]-HeptaMangan

EC FERTILISER | Manganese complex with heptagluconic acid

ρ 1,3 | pH 6,0 - 7,5

Nutrients (w/w)	g/l		
5 % Water-soluble manganese (Mn)	65		
Cross with sutriant deficiency will be more successfully assignt discass, and chiefle stress. Foliar fartilization with			

Crops with nutrient deficiency will be more susceptible against diseases and abiotic stress. Foliar fertilization with macro-and micro-elements will ensure an optimized plant nutrition.

Crop	Aim/Problem	Recommendation	Time
In all crops	To provide manganese, leaf quality, yield, water balance, photosynthesis rate, reduction in radiation stress (antioxidant).	Numerous applications of 2 - 4 l/ha (in at least 200 l water. Upon application with backpack sprayer 0.5 - 1%.)	When required
In all crops	To provide manganese, leaf quality, yield, water balance, photosynthesis rate, reduction in radiation stress (antioxidant).	Fertigation	Ask your consultant
In all crops	Seed dressing with nutrients for improved early growth development and vitality	0,2 - 0,4 l/dt	For seed/plant seed dressing
Cereals	N efficiency, vitality, tillering, stem stability, winter hardiness	2 - 4 times 2 - 4 l/ha	From 3-leaf-stage
Pasture land	N efficiency, vitality, tillering, stem stability, winter hardiness	2 - 5 times 2 - 4 l/ha	During the vegetation period
Potatoes	N efficiency, vitality, fast early growth development, skin quality	3 l/ha	As seed dressing
Potatoes	N efficiency, vitality, skin quality	2 - 4 times 2 - 4 l/ha	From 6-leaf stage
Legumes	N efficiency, increased vitality (e. g. in cold conditions)	2 - 3 times 2 - 4 l/ha	From 6-leaf stage
Maize	N efficiency, increased vitality (e. g. in cold conditions)	1 - 2 times 2 - 4 l/ha	From 4-leaf stage
Oilseed rape	N efficiency, vitality, oil yield, winter hardiness	2 - 3 times 2 - 4 l/ha	From 4-leaf stage
Sunflowers	N efficiency, vitality, oil yield	2 - 4 l/ha	From 4-leaf stage
Sugar beet	N efficiency, increased vitality (e. g. in cold conditions)	1 - 3 times 2 - 4 l/ha	From 6-leaf stage
Strawberries	N efficiency, vitality, winter hardiness	2 - 4 times 2 l/ha	From green buds
Pome fruit	N efficiency, increased vitality (e.g. in cold conditions), fruit colouration	2 - 4 times 2 l/ha	Red bud until harvest
Stone fruit	N efficiency, increased vitality (e. g. in cold conditions)	2 - 3 times 2 l/ha	From fruit set
Soft fruit	N efficiency, increased vitality (e. g. in cold conditions)	2 - 3 times 2 l/ha	Start of shoot growth
Dessert grapes	N efficiency, increased vitality (e. g. in cold conditions)	2 - 3 times 2 l/ha	Inflorescences visible

You can find more information on the hotline: +49 (0) 63 28-9 84 94-80 or on our website www.lebosol.de



Dünger GmbH | Wiesengasse 28 | D-67471 Elmstein | Tel. +49 (0) 63 28-9 84 94-0 | info@lebosol.de | www.lebosol.de

Сгор	Aim/Problem	Recommendation	Time
Citrus fruits	N efficiency, increased vitality (e. g. in cold conditions)	2 - 3 times 3 l/ha	From white bud to harvest
Wine grapes	N efficiency, increased vitality (e. g. in cold conditions)	2 - 3 times 2 l/ha	Inflorescences visible
Medicinal plants, scented plants and spice plants	Leaf quality, N efficiency, increased vitality (e.g. in cold conditions), oil yield	2 - 4 times 2 - 4 l/ha	Once sufficient leaf mass has developed
General vegetables	N efficiency, increased vitality (e. g. in cold conditions)	2 - 4 times 2 - 4 l/ha	Once sufficient leaf mass has developed
Hops	N efficiency, vitality, winter hardiness	1 - 4 times 3 l/ha	From 0.5 m growth height
Tobacco	N efficiency, increased vitality (e. g. in cold conditions)	1 - 3 times 3 l/ha	From 4-leaf stage
Christmas trees	N efficiency, vitality, needle quality, winter hardiness	2 - 3 times 2 l/ha	From budding
Ornamental plants	Leaf quality, vitality	2 times 2 l/ha	Once sufficient leaf mass has developed
Greens	N efficiency, vitality, tillering, stem stability, winter hardiness	2 - 5 times 2 - 4 l/ha	During the vegetation period
Nuts	N efficiency, increased vitality (e.g. in cold conditions), skin quality	2 - 5 times 3 l/ha	From swelling buds
Cotton	N efficiency, increased vitality (e.g. in cold conditions), winter hardiness	2 - 3 times 2 - 4 l/ha	From 4-leaf stage
Rice	N efficiency, vitality, tillering, stem stability	2 - 5 times 2 - 4 l/ha	From 3-leaf-stage

You can find more information on the hotline: +49 (0) 63 28-9 84 94-80 or on our website www.lebosol.de



Dünger GmbH | Wiesengasse 28 | D-67471 Elmstein | Tel. +49 (0) 63 28-9 84 94-0 | info@lebosol.de | www.lebosol.de